arXiv:1111.3087v1 [math.DG] 14 Nov 2011

THE ALEXANDROV PROBLEM IN A QUOTIENT SPACE OF $\mathbb{H}^2 \times \mathbb{R}$

Ana Maria Menezes

Abstract

In this paper we prove an Alexandrov type theorem for a quotient space of $\mathbb{H}^2 \times \mathbb{R}$. More precisely we classify the compact embedded surfaces with constant mean curvature in the quotient of $\mathbb{H}^2 \times \mathbb{R}$ by a subgroup of isometries generated by a parabolic translation along horocycles of $\mathbb{H}^2$ and a vertical translation. Moreover, we construct some examples of periodic minimal surfaces in $\mathbb{H}^2 \times \mathbb{R}$.

1 Introduction

Alexandrov's famous theorem, proved in 1962, is a theorem of classification. It tells us that the only compact embedded constant mean curvature hypersurface in $\mathbb{R}^n, \mathbb{H}^n$ and $\mathbb{S}^n_+$ is the round sphere. Since then, some people have proved an Alexandrov type theorem in other spaces.

For instance, in [8] W.T. Hsiang and W.Y. Hsiang proved that a compact embedded constant mean curvature surface in $\mathbb{H}^2 \times \mathbb{R}$ or in $\mathbb{S}^2_+ \times \mathbb{R}$ is a rotational sphere. They used the Alexandrov reflection method with vertical planes in order to prove that for any horizontal direction, there is a vertical plane of symmetry of the surface, orthogonal to that direction. To apply the Alexandrov reflection method we need to start with a vertical plane orthogonal to a given direction that does not intersect the surface, and in $\mathbb{S}^2 \times \mathbb{R}$, this fact is guaranted by the hypothesis that the surface is contained in a hemisphere times $\mathbb{R}$. We remark that in $\mathbb{S}^2 \times \mathbb{R}$, we know there are embedded rotational constant mean curvature tori, but the Alexandrov problem is not completely solved, as in $\mathrm{Nil}_3, \widetilde{\mathrm{PSL}_2}(\mathbb{R})$ and Berger spheres too.

In Sol_3, Rosenberg proved an embedded compact constant mean curvature surface is a sphere [1].

Recently, Mazet, Rodríguez and Rosenberg [4] considered the quotient of $\mathbb{H}^2 \times \mathbb{R}$ by the discrete group of isometries of $\mathbb{H}^2 \times \mathbb{R}$ generated by a horizontal translation along geodesics of $\mathbb{H}^2$ and a vertical translation. In this space they proved an Alexandrov type theorem, i.e., they classified the compact embedded constant mean curvature surfaces in this quotient space. Moreover, they contructed examples of periodic minimal surfaces in $\mathbb{H}^2 \times \mathbb{R}$, where by periodic we mean a surface which is invariant by a non-trivial discrete group of isometries of $\mathbb{H}^2 \times \mathbb{R}$.

In this paper we also consider periodic surfaces in $\mathbb{H}^2 \times \mathbb{R}$. The discrete groups of isometries of $\mathbb{H}^2 \times \mathbb{R}$ we consider are generated by a horizontal translation ψ along horocycles $c(s)$ of $\mathbb{H}^2$ and/or a vertical translation $T(h)$ by some $h > 0$. In the case the group is the $\mathbb{Z}^2$ subgroup of the isometry group generated by ψ and $T(h)$, the quotient space $\mathcal{M} = \mathbb{H}^2 \times \mathbb{R}/[\psi, T(h)]$ is diffeomorphic but not isometric to $\mathbb{T}^2 \times \mathbb{R}$, where $\mathbb{T}^2$ is the 2-torus. Moreover, $\mathcal{M}$ is foliated by the family of tori $\mathbb{T}(s) = c(s) \times \mathbb{R}/[\psi, T(h)]$ which are intrinsically flat and have constant mean curvature $\frac{1}{2}$.

This paper is organized as follows. In section 2 we fix some notation. In Section 3 we classify the compact embedded constant mean curvature surfaces in such a $\mathcal{M}$, i.e., we prove an Alexandrov type theorem for doubly periodic H-surfaces (see Theorem 1). In section 4 we construct some examples of periodic minimal surfaces in $\mathbb{H}^2 \times \mathbb{R}$. In section 5 we prove a multi-valued Rado theorem for small perturbations of the helicoid (see Theorem 2).

Acknowledgements. This work is part of the author's Ph.D. thesis at IMPA. The author would like to express her sincere gratitude to her advisor Prof. Harold Rosenberg for his constant encouragement and guidance throughout the preparation of this work. The author was financially suported by CNPq-Brazil.

2 Preliminaries

Throughout this paper, the Poincaré disk model is used for the hyperbolic plane, i.e.

$$\mathbb{H}^2 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 1\}$$

with the hyperbolic metric $g_{-1} = \frac{4}{(1-x^2-y^2)^2} g_0$, where g_0 is the Euclidean metric in $\mathbb{R}^2$. In this model, the asymptotic boundary $\partial_\infty \mathbb{H}^2$ of $\mathbb{H}^2$ is identified with the unit circle. Consequently, any point in the closed unit disk is viewed as either a point in $\mathbb{H}^2$ or a point in $\partial_\infty \mathbb{H}^2$. We denote by $\mathbf{0}$ the origin of $\mathbb{H}^2$.

In $\mathbb{H}^2$ we denote by γ_0, γ_1 the geodesic lines $\{x = 0\}, \{y = 0\}$, respectively. For $j = 0, 1$, we denote by Y_j the Killing vector field whose flow $(\phi_l)_{l \in (-1,1)}$ is the hyperbolic translation along γ_j with $\phi_l(\mathbf{0}) = (l \sin \pi j, l \cos \pi j)$ and has $(\sin \pi j, \cos \pi j)$ as attractive point at infinity. We call $(\phi_l)_{l \in (-1,1)}$ the flow of Y_j, even though the family $(\phi_l)_{l \in (-1,1)}$ is not parameterized at the right speed.

We write t for the height coordinate in $\mathbb{H}^2 \times \mathbb{R}$ and denote by $\pi : \mathbb{H}^2 \times \mathbb{R} \to \mathbb{H}^2$ the vertical projection. In what follows, we will often identify the hyperbolic plane $\mathbb{H}^2$ with the horizontal slice $\{t = 0\}$ of $\mathbb{H}^2 \times \mathbb{R}$. The vector fields $Y_j, j = 0, 1$ and their flows naturally extend to horizontal vector fields and their flows in $\mathbb{H}^2 \times \mathbb{R}$.

Denote by $(c(s))_{s\in\mathbb{R}}$ the family of horocycles in $\mathbb{H}^2$ tangent to $(1,0)$ at infinity. Given two points $p, q \in c(s)$, we denote by ψ the parabolic translation along $c(s)$ such that $\psi(p) = q$.



We write $\overline{pq}$ to denote the geodesic arc between the two points p, q of $\mathbb{H}^2 \times \mathbb{R}$.

3 The Alexandrov problem for doubly periodic constant mean curvature surfaces

Take two points $p, q \in c(s)$, for some s, and let ψ be the parabolic translation along $c(s)$ such that $\psi(p) = q$. We have $\psi(c(s)) = c(s)$ for all s. Consider G the $\mathbb{Z}^2$ subgroup of Isom$(\mathbb{H}^2 \times \mathbb{R})$ generated by ψ and $T(h)$, for some positive h. We denote by $\mathcal{M}$ the quotient of $\mathbb{H}^2 \times \mathbb{R}$ by G. The manifold $\mathcal{M}$ is diffeomorphic to $\mathbb{T}^2 \times \mathbb{R}$. Moreover, $\mathcal{M}$ is foliated by the family of tori $\mathbb{T}(s) = (c(s) \times \mathbb{R})/G$, $s \in \mathbb{R}$, which are intrinsically flat and have constant mean cuvature $\frac{1}{2}$. Therefore, the tori $\mathbb{T}(s)$ are examples of compact embedded constant mean curvature surfaces in $\mathcal{M}$.

We have the following answer to the Alexandrov problem in $\mathcal{M}$.

Theorem 1. *Let* $\Sigma \subset \mathcal{M}$ *be a compact embedded surface with constant mean curvature* H*. Then* $H \geq \frac{1}{2}$*. Moreover,*

1. If $H = \frac{1}{2}$*, then* Σ *is a torus* $\mathbb{T}(s)$*, for some* s; *and*

2. If $H > \frac{1}{2}$, Σ *is either the quotient of a rotational sphere, or the quotient of a vertical unduloid (in particular, a vertical cylinder over a circle).*

Proof. Let Σ be a compact embedded surface in $\mathcal{M}$ with constant mean curvature H. As Σ is compact, there exist $s_0 \leq s_1 \in \mathbb{R}$ such that Σ is between $\mathbb{T}(s_1)$ and $\mathbb{T}(s_0)$ and it is tangent to $\mathbb{T}(s_0), \mathbb{T}(s_1)$ at points p, q, respectively, as illustrated in the figure below.

For $s > s_1, \mathbb{T}(s)$ does not touch Σ and Σ stays in the mean convex region bounded by $\mathbb{T}(s_1)$. By comparison at q, we conclude that $H \geq \frac{1}{2}$. If $H = \frac{1}{2}$, then by the maximum principle Σ is the torus $\mathbb{T}(s_1)$, and we have already proved the first part of the thereom.

To proof the last part, consider a connected component $\widetilde{\Sigma}$ of the lift of Σ to $\widetilde{\mathcal{M}} = \mathbb{H}^2 \times \mathbb{R}/[\mathbb{T}(h)] = \mathbb{H}^2 \times \mathbb{S}^1$. Denote by $\tilde{c}(s)$ the surface $c(s) \times \mathbb{S}^1$. Note that $\tilde{c}(s)$ is the lift of $\mathbb{T}(s)$ to $\widetilde{\mathcal{M}}$.



Claim. $\widetilde{\Sigma}$ separates $\widetilde{\mathcal{M}} = \mathbb{H}^2 \times \mathbb{S}^1$.

In fact, if it is not the case, we can consider a geodesic arc $\gamma : (-\epsilon, \epsilon) \to \widetilde{\mathcal{M}}$ such that $\gamma(0) \in \widetilde{\Sigma}, \gamma'(0) \in T\widetilde{\Sigma}^{\perp}$ and we can join the points $\gamma(-\epsilon), \gamma(\epsilon)$ by a curve that does not intersect $\widetilde{\Sigma}$, hence we obtain a jordan curve whose intersection number with $\widetilde{\Sigma}$ is 1 modulo 2. Note that each point of $\widetilde{\Sigma}$ has bounded distance to $\tilde{c}(s_0)$. Since we can homotop the jordan curve such that its points are an arbitrary distance from $\tilde{c}(s_0)$, we conclude that this jordan curve does not intersect $\widetilde{\Sigma}$ anymore, contradicting the fact that its intersection number is 1 modulo 2. Thus the claim is proved.

Let us call A the mean convex component of $\widetilde{\mathcal{M}} \setminus \widetilde{\Sigma}$ with boundary $\widetilde{\Sigma}$ and B the other component, hence $\widetilde{\mathcal{M}} \setminus \widetilde{\Sigma} = A \cup B$. Also, let $\tilde{p}, \tilde{q} \in \widetilde{\mathcal{M}}$ whose projections in $\mathcal{M}$ are p, q respectively.

Let γ be a geodesic in $\mathbb{H}^2$ that limits to $(1,0) \in \partial_\infty \mathbb{H}^2, \gamma(-\infty) = (1,0)$ (the point where the horocycles $c(s)$ are centered) and consider $(l_t)_{t \in \mathbb{R}}$ the family of geodesics in $\mathbb{H}^2$ orthogonal to γ. Denote by $P(t)$ the totally geodesic vertical annulus $l_t \times \mathbb{S}^1$ of $\widetilde{\mathcal{M}} = \mathbb{H}^2 \times \mathbb{S}^1$. Since $\widetilde{\Sigma}$ is a lift of the compact surface Σ, it stays in the region between $\tilde{c}(s_1)$ and $\tilde{c}(s_0)$, and the distance from any point of $\widetilde{\Sigma}$ to $\tilde{c}(s_1)$ and to $\tilde{c}(s_0)$ is uniformly bounded.



For t large, $P(t)$ is contained in B and disjoint from $\tilde{c}(s_1)$. Moreover, $P(t)$ has points at an arbitrary distance from $\tilde{c}(s_1)$. It follows that $P(t) \cap \widetilde{\Sigma}$ is compact for all t, and is empty if $|t|$ is large enough. Then start with t close to infinity and let t decrease until a first contact point between $\widetilde{\Sigma}$ and $P(t_0)$. By the maximum principle, we know that the mean curvature vector of $\widetilde{\Sigma}$ does not point into $\bigcup_{t \geq t_0} P(t)$.

Let us suppose $\widetilde{\Sigma} \cap \gamma$ has at least two points. Continuing to decrease t and start Alexandrov reflection procedure for $\widetilde{\Sigma}$ and the family of vertical totally geodesic annuli $P(t)$, we get a first contact point between the reflected part of $\widetilde{\Sigma}$ and $\widetilde{\Sigma}$, for some $t_1 \in \mathbb{R}$. Then $\widetilde{\Sigma}$ is symmetric with respect to $P(t_1)$. But $\widetilde{\Sigma} \cap \left(\bigcup_{t_1 \leq t \leq t_0} P(t) \right)$ is compact, and then $\widetilde{\Sigma}$ is compact. Hence, given any horizontal geodesic β that limits to $(1,0)$ (or to any other point in $(1,0) \times \mathbb{S}^1$) we can apply the Alexandrov procedure with the family of totally geodesic vertical annulus $Q(t) = \tilde{l}_t \times \mathbb{S}^1$, where $(\tilde{l}_t)_{t \in \mathbb{R}}$ is the family of horizontal geodesics orthogonal to β, and we obtain a symmetry plane.

Hence we have shown that if some geodesic that limits to $(1,0)$ intersects $\widetilde{\Sigma}$ in two or more points, then $\widetilde{\Sigma}$ lifts to a rotational cylindricaly bounded surface $\bar{\Sigma}$ in $\mathbb{H}^2 \times \mathbb{R}$. If $\bar{\Sigma}$ is non compact then $\bar{\Sigma}$ is a Delauney surface and if $\bar{\Sigma}$ is compact, by the Hsiang-Hsiang's theorem, we know that $\bar{\Sigma}$ is a rotational sphere.

Now, let us suppose every geodesic that limits to $(1,0)$ intersects $\widetilde{\Sigma}$ in at most one point. In particular, the geodesic β that limits to $(1,0)$ and passes through $\tilde{p} \in \tilde{c}(s_0)$, intersects $\widetilde{\Sigma}$ only at $\tilde{p}$. Write β^- to denote the arc of β between $\tilde{p}$ and $\beta(+\infty)$.



As $\beta \cap \widetilde{\Sigma} = \{\tilde{p}\}$, we have $\beta^- \cap \widetilde{\Sigma} = \emptyset$ and then $\beta^- \subset B$.

Hence, at the point $\tilde{p} \in \widetilde{\Sigma} \cap \tilde{c}(s_0)$, the mean curvature vectors of $\widetilde{\Sigma}$ and $\tilde{c}(s_0)$ point in the same direction and $\widetilde{\Sigma}$ is on one side of $\tilde{c}(s_0)$, then by comparison we get $H \leq \frac{1}{2}$. But we already know that $H \geq \frac{1}{2}$. Hence, $H = \frac{1}{2}$ and $\widetilde{\Sigma} = \tilde{c}(s_0)$ by the maximum principle. Therefore, $\Sigma = \mathbb{T}(s_0)$.

□

4 Periodic minimal surfaces

In this section we are interested in constructing minimal surfaces which are invariant by a subgroup of $\mathrm{Isom}(\mathbb{H}^2 \times \mathbb{R})$ isomorphic to $\mathbb{Z}^2$ or generated by a vertical translation. In fact, we only consider subgroups generated by a parabolic translation ψ along a horocycle and/or a vertical translation $T(h)$, for some $h > 0$.

4.1 Doubly periodic minimal surface

Denote by γ_0, γ_1 the geodesic lines in $\mathbb{H}^2$, $\{x = 0\}, \{y = 0\}$, respectively. Let c be a horocycle orthogonal to γ_1, and consider $p, q \in c$ equidistant points to γ_1. Take α, β geodesics which limit to $(1, 0)$ and pass through p, q, respectively. We have $p = \alpha(t_0), q = \beta(t_0)$, for some t_0. Consider a point $A = \gamma_1(t_0)$. Parametrize so that when $t_0 \to \infty$, we have $\alpha(t_0) \to B$, $\beta(t_0) \to D$ and $\gamma_1(t_0) \to (1, 0)$.



We fix $h > \pi$ and consider the following Jordan curve:

$$\begin{aligned}\Gamma_{t_0} \quad = \quad & \overline{(p,0),(A,0)} \cup \overline{(p,0),(p,h)} \cup \overline{(q,0),(A,0)} \\ & \cup \overline{(q,0),(q,h)} \cup \overline{(p,h),(A,h)} \cup \overline{(q,h),(A,h)}.\end{aligned}$$



Figure 1: Γ_{t_0}

We consider a least area embedded minimal disk Σ_{t_0} with boundary Γ_{t_0}. As Γ_{t_0} is tranversal to the Killing field Y_1, we can show using the maximum principle that Σ_{t_0} is a Y_1-Killing graph with convex boundary, in particular, Σ_{t_0} is stable and unique (see Lemma 2.1 in [3]). This gives uniform curvature estimates far from the boundary. Following Appendix C in [4], we can show $\Sigma_{t_0} \cap \{0 \leq t \leq \frac{h}{2}\}$

is a vertical graph and then it can be extended by symmetry with respect to $\overline{(p,0),(A,0)}$ to a larger vertical graph. Thus we also obtain uniform curvature estimates in a neighborhood of $\overline{(p,0),(A,0)}$. This is also true for the three other horizontal geodesic arcs in Γ_{t_0}.

Observe that, by the maximum principle, for any t_0, Σ_{t_0} stays in the half-space determined by $\overline{BD} \times \mathbb{R}$ that contains Γ_{t_0}.

Since $h > \pi$, we can consider the minimal surface $S_h \subset \mathbb{H}^2 \times (0,h)$ which is a vertical bigraph with respect to $\{t = \frac{h}{2}\}$, it is invariant by tranlations along the horizontal geodesic γ_0 and whose boundary is $(\tau \times \{0\}) \cup \overline{(0,1,0)(0,1,h)}$ $\cup (\tau \times \{h\}) \cup \overline{(0,-1,0)(0,-1,h)}$, where $\tau = \partial_\infty \mathbb{H}^2 \cap \{x > 0\}$. For more details about the surface S_h, see [4, 5, 6].

Let $(\phi_l)_{l \in \mathbb{R}}$ be the flow of the Killing vector field Y_1. For l close to 1, the translated surface $\phi_l(S_h)$ does not meet Σ_{t_0}. The surface Σ_{t_0} is contained between $\phi_l(S_h)$ and $\overline{BD} \times \mathbb{R}$. When $t_0 \to +\infty$, $\Gamma_{t_0} \to \Gamma$, where

$$\begin{aligned} \Gamma \;=\; & (\alpha \times \{0\}) \cup (\beta \times \{0\}) \cup (\alpha \times \{h\}) \cup \\ & (\beta \times \{h\}) \cup \overline{(D,0)(D,h)} \cup \overline{(B,0)(B,h)}. \end{aligned}$$

Using the maximum principle, we have that for all $t > t_0$, Σ_t is contained between $\phi_l(S_h)$ and $\overline{BD} \times \mathbb{R}$. Therefore, the surface Σ_{t_0} converges to a minimal surface Σ, where Σ lies in the region of $\{0 \leq t \leq h\}$ bounded by $\alpha \times \mathbb{R}$, $\beta \times \mathbb{R}$, $\overline{BD} \times \mathbb{R}$ and $\phi_l(S_h)$; $\partial\Sigma = \Gamma$ and $\Sigma \setminus \partial\Sigma$ is a vertical bigraph with respect to $\{t = \frac{h}{2}\}$.

Hence, the surface obtained by reflection in all horizontal boundary geo- desics of Σ is invariant by ψ^2 and $T(2h)$. Moreover, this surface in the quotient space $\mathbb{H}^2 \times \mathbb{R} / [\psi^2, T(2h)]$ is topologically a sphere minus three points.

4.2 Vertically periodic minimal surfaces

In this section we will construct some examples of minimal surfaces invariant by a vertical translation.

Take α any geodesic in $\mathbb{H}^2 \times \{0\}$. For $h > \pi$, consider the asymptotic vertical segment $\alpha(+\infty) \times [0, 2h]$, and a point $p \in \partial_\infty \mathbb{H}^2$. For some small $\epsilon > 0$, consider the vertical segment joining $p \times \{\epsilon\}$ and $p \times \{h + \epsilon\}$. Now, connect $p \times \{\epsilon\}$ to $\alpha(+\infty) \times \{0\}$ and $p \times \{h + \epsilon\}$ to $\alpha(+\infty) \times \{2h\}$ by curves in $\partial_\infty \mathbb{H}^2 \times \mathbb{R}$, whose tangent vectors never point in the horizontal or vertical directions, and so that the resulting curve Γ is differentiable. Also, consider the horizontal geodesic β connecting p to $\alpha(-\infty)$.

Write α_t to denote $\alpha(t) \times [0, 2h]$, and for each t consider a curve Γ_t with endpoints $\alpha(t) \times \{0\}$ and $\alpha(t) \times \{2h\}$, contained in the region R bounded by $\alpha \times \mathbb{R}, \beta \times \mathbb{R}, \mathbb{H}^2 \times \{0\}$ and $\mathbb{H}^2 \times \{2h\}$, such that its tangent vector never points in the horizontal direction and $\Gamma_t \to \Gamma$ when $t \to +\infty$.



Fix t. For each n, let Σ_t^n be the solution to the plateau problem with boundary $\Gamma_t \cup \alpha([t-n,t]) \times \{0\} \cup \alpha([t-n,t]) \times \{2h\} \cup \alpha_{t-n}$. Given γ any geodesic orthogonal to α, if we consider Y the Killing field which is the translation along γ, we have that $\partial\Sigma_t^n$ is transverse to Y and then, using the maximum principle, we conclude that Σ_t^n is a Y-Killing graph, in particular is unique and stable. Hence, by the Main Theorem in [7], we have uniform curvature estimates. Therefore, when n goes to infinity, Σ_t^n converges to a minimal surface Σ_t with $\alpha(-\infty) \times [0,2h] \subset \partial_\infty \Sigma_t$. We know the surface Σ_t^n is contained between $\alpha \times \mathbb{R}$ and $\beta \times \mathbb{R}$, then we have $\partial_\infty \Sigma_t = \alpha(-\infty) \times [0,2h]$ necessarily.

For each t, $\Sigma_t \subset R$ and $\Gamma_t \to \Gamma$, when $t \to +\infty$. We will show that a subsequence of Σ_t converges to a minimal surface Σ with $\Gamma \subset \partial_\infty \Sigma$. We use suitable translations of the surface S_h, mentioned before, as a barrier. By our choice of Γ, for each $q \in \Gamma$, we can consider two translations of S_h that pass through q so that one of them has asymptotic boundary under Γ and the other one has asymptotic boundary above Γ. Hence, the envelope of the union of all these translated surfaces S_h forms a barrier to Σ_t, for all t, and then we conclude that $\partial_\infty \Sigma = \Gamma \cup \alpha(-\infty) \times [0,2h]$.

Therefore, the surface obtained by reflection in all horizontal boundary geodesics of Σ is invariant by $T(4h)$.

Now we construct another example of vertically invariant minimal surface. Consider two geodesics α, β in $\mathbb{H}^2$ that limit to a same point at infinity, say $\alpha(-\infty) = (1,0) = \beta(-\infty)$. For $h > \pi$, consider a curve Γ contained in the asymptotic boundary of $\mathbb{H}^2 \times \mathbb{R}$, connecting $\alpha(+\infty) \times \{2h\}$ to $\beta(+\infty) \times \{0\}$ such that its tangent vector never points in the horizontal or vertical directions. Also, take a point $p \in \partial_\infty \mathbb{H}^2$ in the halfspace determined by $\beta \times \mathbb{R}$ that does not contain α.

For some small $\epsilon > 0$, consider the vertical segment joining $p \times \{\epsilon\}$ and $p \times \{h+\epsilon\}$. Now, connect $p \times \{\epsilon\}$ to $(1,0) \times \{0\}$ and $p \times \{h+\epsilon\}$ to $(1,0) \times \{2h\}$ by curves in $\partial_\infty \mathbb{H}^2 \times \mathbb{R}$, whose tangent vectors never point in the horizontal or vertical directions, and such that the resulting curve $\widehat{\Gamma}$ is differentiable.

For each t, s, consider two curves $\widehat{\Gamma}_t$ and Γ_s contained in the region R bounded by $\alpha \times \mathbb{R}$, $\beta \times \mathbb{R}$, $\mathbb{H}^2 \times \{0\}$ and $\mathbb{H}^2 \times \{2h\}$, joining the point $\alpha(t) \times \{2h\}$ to $\beta(t) \times \{0\}$

and $\alpha(s) \times \{2h\}$ to $\beta(s) \times \{0\}$, respectively, such that their vector tangents are never horizontal and $\Gamma_s \to \Gamma$ when $s \to +\infty$ and $\widehat{\Gamma}_t \to \widehat{\Gamma}$ when $t \to -\infty$.



Fix t. For each $s > t$, let Σ_t^s be the solution to the plateau problem with boundary $\Gamma_s \cup \alpha([t,s]) \times \{2h\} \cup \widehat{\Gamma}_t \cup \beta([t,s]) \times \{0\}$. The surface Σ_t^s is contained in the region R bounded $\alpha \times \mathbb{R}, \beta \times \mathbb{R}, \mathbb{H}^2 \times \{0\}$ and $\mathbb{H}^2 \times \{2h\}$. When s goes to infinity, a subsequence of Σ_t^s converges to a minimal surface Σ_t with $\Gamma \subset \partial_\infty \Sigma_t$. We use suitable translations of the surface S_h, mencioned before, as a barrier. By our choice of Γ, for each $q \in \Gamma$, we can consider two translations of S_h that pass through q such that one of them has asymptotic boundary above Γ, the other one under Γ and their intersection with Γ is just the point q considered. Hence, the envelope of the union of all these translated surfaces S_h forms a barrier to Σ_t^s for all s, and then we conclude that $\partial_\infty \Sigma_t = \Gamma$.

Now, let decrease t. As $\widehat{\Gamma}_t \to \widehat{\Gamma}$, a subsequence of Σ_t converges to a minimal surface Σ with $\widehat{\Gamma} \subset \partial_\infty \Sigma$, and using the same idea as in the previous section we can show $\widehat{\Gamma} = \partial_\infty \Sigma$.

The surface obtained by reflection in all horizontal boundary geodesics of Σ is a minimal surface invariant by $T(4h)$.

5 A perturbation of the Helicoid

The aim of this section is to proof a multi-valued Rado theorem for small perturbation of the helicoid. Recall that we denote by $t : \mathbb{H}^2 \times \mathbb{R} \to \mathbb{R}$ the horizontal projection.

Consider Y the Killing field whose flow $\phi_\theta, \theta \in [0, 2\pi)$, is the rotation by angle θ around the z-axis. For some $0 < c < 1$, denote $D = \{(x,y) \in \mathbb{H}^2; x^2 + y^2 \leq c\}$. Take a helice h_0 of constant pitch cointained in a bounded slab, say $S = \{p \in \mathbb{H}^2 \times \mathbb{R}; 0 \leq t(p) \leq d\}$, in such a way that the vertical projection of h_0 over $H^2 \times \{0\}$ is ∂D. Consider h a small perturbation of this helice with fixed endpoints such that h is transversal to Y. Now take the two horizontal geodesics joining the endpoints of h to the z-axis. Call Γ the Jordan curve which is the union of h, the two horizontal geodesics and a part of the z-axis, as illustrated in the figure below.



Let H be the solution to the plateau problem with boundary Γ. Then, in particular, H is a compact minimal disk with $\partial H = \Gamma$.

Theorem 2 (A multi-valued Rado Theorem)**.** *Under the assumptions above, H is the unique compact minimal disk with boundary Γ.*

Proof. Let us denote $\Gamma_\theta = \phi_\theta(\Gamma)$ and $H_\theta = \phi_\theta(H)$, hence H_θ is a minimal disk with $\partial H_\theta = \Gamma_\theta$. As h is transversal to Y, $(H_\theta)_{\theta \in [0,2\pi)}$ gives us a foliation of $R = \pi^{-1}(D) \cap S$.

Let $M \neq H$ be another compact minimal disk with boundary Γ. We will analyse the intersection between M and each H_θ.

Firstly, observe that $M \cap H_\theta \neq \emptyset$ for all θ and by the maximum principle $M \subset R$.

Fix θ_0. Given $q \in H_{\theta_0} \cap M$, either $q \in \text{int}M$ or $q \in \Gamma = \partial M$.

Suppose $q \in \text{int}M$. If the intersection is transversal at q, then in a neighborhood of q we have $H_{\theta_0} \cap M$ is a simple curve passing through q. On the other hand, if M is tangent to H_{θ_0} at q, in a neighborhood of q we have $H_{\theta_0} \cap M$ is $2k$ curves passing through q, making equal angles at q. If we let θ_0 vary a little, we see in M a foliation as in the figure below.



$T_qM \neq T_qH_{\theta_0}$

or



$T_qM = T_qH_{\theta_0}$

Now suppose $q \in \Gamma$. If $q \in \Gamma \cap \{z\text{-axis}\}$, we may obtain a θ such that $T_qM = T_qH_\theta$. As $M \neq H$, M is not on one side of H_θ and we see k curves starting at q contained in $M \cap H_\theta$. Moreover, if we vary θ a little, we obtain the following foliation in a neighborhood of q (see Figure 2).

On the other hand, if $q \in \Gamma \setminus \{z\text{-axis}\}$, since for $\theta \neq 0$, $\Gamma_\theta \cap \Gamma \setminus \{z\text{-axis}\} = \emptyset$, then $\theta_0 = 0$, and we can not have $M \cap H_\theta$ homeomorphic to a semicircle in a



Figure 2: $q \in \Gamma \cap \{z\text{-axis}\}$

neighborhood of q. (See Figure 3.) Since then M is on one side of H_θ at q which contradicts the boundary maximum principle.



Figure 3:

Thus, for θ close to 0, in a neighborhood of q we have that $M \cap H_\theta$ is as in Figure 4.



Figure 4: $q \in \Gamma \setminus \{z\text{-axis}\}$

Consider two copies of M and glue them identifying equal in the boundary.

As M is a disk, when we glue these two copies of M we obtain a sphere with a foliation with singularities, each one with negative index. But this is impossible. Therefore, there is no other minimal disk with boundary Γ besides H.

□

Consider h_1 a curve in $\partial_\infty \mathbb{H}^2 \times \mathbb{R}$ which is a small perturbation of a helice and the distance between two points in h_1 in a same vertical line is greater than π and call Γ_1 the Jordan curve which is the union of h_1, the two horizontal geodesics joining the endpoints of h_1 to the z-axis and a part of the z-axis. Denote by h_t a perturbation of a helice over the circle $\{(x,y) \in \mathbb{H}^2, x^2 + y^2 = t\}$ as before. Call Γ_t the Jordan curve which is the union of h_t, the two horizontal geodesics joining the endpoints of h_t to the z-axis and a part of the z-axis. We choose h_t so that $\Gamma_t \to \Gamma_1$ when $t \to 1$. Denote by H_t the minimal disk with boundary Γ_t. We have H_t is stable and unique. Thus, as in the previous section we can use the barriers $S_a, a > \pi$, to prove that when $t \to 1$, the limit surface H_1 exists and has $\partial H_1 = \Gamma_1$.

References

[1] B. Daniel and P. Mira. Existence and uniqueness of constant mean curvature spheres in Sol_3. Preprint, 2008, arXiv:0812.3059.

[2] B. Nelli and H. Rosenberg. Minimal surfaces in $\mathbb{H}^2 \times \mathbb{R}$. Bull. Braz. Math. Soc., 33(2), 263-292, 2002.

[3] B. Nelli and H. Rosenberg. Simply connected constant mean curvature surfaces in $\mathbb{H}^2 \times \mathbb{R}$. Mich. Math. J. 54(3), 537-544, 2006.

[4] L. Mazet, M. Magdalena Rodríguez and H. Rosenberg. Periodic constant mean curvature surfaces in $\mathbb{H}^2 \times \mathbb{R}$. Preprint, 2011, arXiv:1106.5900.

[5] L. Mazet, M. Magdalena Rodríguez and H. Rosenberg. The Dirichlet problem for minimal surface equationc - with possible infinite boundary date - over domains in riemannian surface. Proc. London Math. Soc. 102(6), 985-1023, 2011.

[6] R. Sa Earp. Parabolic and hyperbolic screw motion surfaces in $\mathbb{H}^2 \times \mathbb{R}$. J. Aust. Math. Soc., 85: 113-143, 2008.

[7] H. Rosenberg, R. Souam and E. Toubiana. General curvature estimates for stable H-surfaces in 3-manifolds and aplications, J. Dif. Geom. 84, 623-648, 2010.

[8] W.T. Hsiang and W.Y. Hsiang. On the uniqueness of isoperimetric solutions and embedded soap bubbles in noncompact symmetric spaces I, Invent.Math. 98(1), 39-58, 1989.

Instituto Nacional de Matemática Pura e Aplicada (IMPA)
Estrada Dona Castorina 110, 22460-320, Rio de Janeiro-RJ, Brazil
Email adress: anamaria@impa.br